

MODIFIED ABOODH HOMOTOPY PERTUBATION METHOD FOR SOLVING NONLINEAR GAS DYNAMIC EQUATION

Olubanwo O. O.¹, Talabi A.T.², Odetunde O. S.¹, Ogunwobi Z. O.¹

¹Department of Mathematics, Olabisi Onabanjo University, Ago-Iwoye, Nigeria.

²Department of Physics, Olabisi Onabanjo University, Ago-Iwoye, Nigeria.

olubanwo.oludapo@oouagoiwoye.edu.ng

ABSTRACT

In this paper, we present a reliable combination of Aboodh Transform and Modified Homotopy perturbation method to solve nonlinear gas dynamic equations. Some problems were solved to demonstrate the capability and reliability of the mixture of Aboodh Transform and Modified Homotopy perturbation method. We have compared the result obtained with the available Laplace Transform New Homotopy Perturbation Method solution and homotopy perturbation method of solution which is found to be exactly the same. The result revealed that the combination of the Aboodh Transform and Modified homotopy perturbation method is practically well appropriate for use in such problems.

Keywords: Aboodh Transform, Homotopy Perturbation Method, Modified Homotopy Perturbation Method, Nonlinear Gas Dynamic Equation, Partial Differential Equation.

1. INTRODUCTION

Nonlinear models has been very useful in describing various phenomena that appear in many areas of scientific fields. We know that except in a limited number of these problems, there are difficulties in obtaining their exact solutions. Thus, many researchers had made several attempts to develop various analytical methods for obtaining solutions which reasonably approximate the exact solutions see Bellman (1964), Cole (1968). In recent years, most nonlinear problems were solved by numerical methods and their improvement led to improvement in analytical methods. Several scientists believe that the combination of numerical and analytical methods will also produce meaningful results. Many new techniques have been widely used to solve nonlinear problems. Example of such is homotopy perturbation method established by He to obtain series solution of nonlinear differential equations (He 1999, 2000a, 2000b, 2003, 2004, 2005, 2006, 2008). The method has the advantages of simplicity and easy execution.

Gas dynamics equation is a mathematical model which is based on the physical laws of conservation, namely: the laws of conservation of mass, conservation of momentum, conservation of energy and many more. The nonlinear equations of ideal gas dynamics are applicable for the three classes of nonlinear waves like shock fronts, rare factions and contact discontinuities. Several methods have been used to solve different classes of gas dynamics equations see Evans and Bulut (2002), Polyanin and Zaitsev (2004), Elizarova (2009), Jafari, Chun, Seifi, and Saeidy (2009), Ames (1965), Rasulov and

Karaguler (2003), Aminikhah and Jamalain (2013), Jafari, Zabihi and Saidy (2008). Recently, Aboodh introduced a new set of integral transform called Aboodh transform and it has been applied to obtain solution of several classes of both linear and nonlinear partial differential equations (Aboodh 2013, 2014, 2015, 2016).

In this paper, we construct the solution of gas dynamic equation by using Modified Aboodh Homotopy Perturbation Method. Analytical approximation to the solution of the nonlinear gas dynamic equation is obtained by using the mixture of Modified Homotopy Perturbation Method and Aboodh Transform. The gas dynamic equation as a nonlinear partial differential model is as follows (Evans and Bulut 2002, Jafari *et al.* 2009, Aminikhah and Jamalain 2013):

$$u_t + uu_x - u(1 - u) = 0 \quad 1$$

where $0 \leq x \leq 1$ and $t > 0$

The result obtained by Modified Aboodh Homotopy Perturbation Method confirm the applicability and effectiveness of the proposed method in solving the nonlinear gas dynamic equation.

2. MODIFIED ABOODH HOMOTOPY PERTUBATION METHOD (MAHPM)

In this section, we propose MAHPM, which is the combination of Aboodh Transform and modified homotopy perturbation methods (MHPM) for solving nonlinear gas dynamic equation. This method is simple and obtain the exact solution of the equations analytically using the initial condition

only. This method provides the solution in a closed form. To illustrate the basic idea of this method, we consider the following nonlinear differential equation (Aminikhah and Jamalian, 2013):

$$A(u) - f(r) = 0, r \in \Omega \tag{2}$$

with the following initial conditions

$$u(0) = \alpha_0, \dot{u}(0) = \alpha_1, \dots, u^{(n-1)}(0) = \alpha_{n-1} \tag{3}$$

where A is a general differential operator and $f(r)$ is a known analytical function. The operator A can be divided into two parts, L and N , where L is a linear and N is a nonlinear operator. Thus, equation (2) can be rewritten as

$$L(u) + N(u) - f(r) = 0 \tag{4}$$

From MHPM [23], we construct a homotopy $U(r, p): \Omega \times [0,1] \rightarrow \mathbb{R}$, which satisfies

$$H(U, p) = (1 - p)[L(U) - u_0] + p[A(U) - f(r)] = 0, p \in [0,1], r \in \Omega \tag{5}$$

$$H(U, p) = L(U) - u_0 + pu_0 + p[N(u) - f(r)] = 0, p \in [0,1], r \in \Omega \tag{6}$$

Where $p \in [0,1]$ is an embedded parameter and u_0 is an initial approximation for the solution of equation (2)

In the case of $p = 0$, then equation (4) becomes

$$H(U, 0) = L(U) - u_0 = 0 \tag{7}$$

In the case of $p = 1$, then equation (5) becomes

$$H(U, 1) = A(U) - f(r) = 0 \tag{8}$$

Now, applying the Aboodh transform to both sides of equation (6), we have

$$A\{L(U) - u_0 + pu_0 + p[N(u) - f(r)]\} = 0 \tag{9}$$

Using the differential property of the aboodh transform we obtain

$$v^n U(x, v) - \frac{U(x,0)}{v^{2-n}} - \frac{U^1(x,0)}{v^{3-n}} - \dots - \frac{U^{(n-1)}(x,0)}{v} = A\{u_0 - pu_0 + p[N(u) - f(r)]\} \tag{10}$$

$$U(x, v) = \frac{1}{v^n} \left\{ \frac{U(x,0)}{v^{2-n}} + \frac{U^1(x,0)}{v^{3-n}} + \dots + \frac{U^{(n-1)}(x,0)}{v} + A\{u_0 - pu_0 + p[N(u) - f(r)]\} \right\} \tag{11}$$

Taking the Inverse Aboodh transform on both sides of equation (11), we have

$$U(x, t) = A^{-1} \left\{ \frac{1}{v^n} \left(\frac{U(x,0)}{v^{2-n}} + \frac{U^1(x,0)}{v^{3-n}} + \dots + \frac{U^{(n-1)}(x,0)}{v} + A\{u_0 - pu_0 + p[N(u) - f(r)]\} \right) \right\} \tag{12}$$

From Homotopy Perturbation Method we considered the embedding parameter p as a small parameter and assume that the solution of equation (12) can be written as a power series of p as:

$$U(x, t) = \sum_{n=0}^{\infty} p^n U_n(x, t) \tag{13}$$

Substituting the equation (12) into (13), we have

$$\sum_{n=0}^{\infty} p^n U_n(x, t) = A^{-1} \left\{ \frac{1}{v^n} \left(\frac{U(x,0)}{v^{2-n}} + \frac{U^1(x,0)}{v^{3-n}} + \dots + \frac{U^{(n-1)}(x,0)}{v} + A\{u_0 - pu_0 + p[N(u) - f(r)]\} \right) \right\} \tag{14}$$

This is the coupling of the Aboodh Transform and the modified homotopy perturbation method. Equating the coefficients of the corresponding powers of p , we have

$$p^0: \quad U_0(x, t) = A^{-1} \left\{ \frac{1}{v^n} \left[\frac{U(x,0)}{v^{2-n}} + \frac{U^1(x,0)}{v^{3-n}} + \dots + \frac{U^{(n-1)}(x,0)}{v} + A\{u_0\} \right] \right\}$$

$$p^1: \quad U_1(x, t) = A^{-1} \left\{ \frac{1}{v^n} \{A\{[N(U_0) - u_0 - f(r)]\}\} \right\}$$

$$p^2: \quad U_2(x, t) = A^{-1} \left\{ \frac{1}{v^n} \{A\{N(U_0, U_1)\}\} \right\}$$

$$p^3: \quad U_3(x, t) = A^{-1} \left\{ \frac{1}{v^n} \{A\{N(U_0, U_1, U_2)\}\} \right\} \text{ and so on}$$

In general, the recursive relation is given by: 15

$$p^m: \quad U_m(x, t) = A^{-1} \left\{ \frac{1}{v^n} \{A\{N(U_0, U_1, U_2, \dots, U_{m-1})\}\} \right\}$$

Assuming that the initial approximate solution of equation (2) has the form $U(0) = u_0 = \alpha_0, U^1(0) = \alpha_1, \dots, U^{(n-1)}(0) = \alpha_{n-1}$. Thus, the exact solution may be obtained as follows:

$$u = \lim_{p \rightarrow 1} U(x, t) = U_0(x, t) + U_1(x, t) + U_2(x, t) + \dots \quad 16$$

3. APPLICATION

In this section, in order to demonstrate the effectiveness and applicability of the Modified Aboodh Homotopy perturbation method for solving gas dynamics equations, we will solve the following examples.

Example 1

Consider the nonlinear gas dynamic equation (Aminikhah and Jamalain 2013)

$$u_t + uu_x - u(1 - u) = 0 \quad 17$$

subject to the constant initial conditions

$$u(x, 0) = ae^{-x} \quad 18$$

By applying the modified homotopy perturbation method, we construct the following homotopy:

$$H(U, p) = U_t - u_0 + p[u_0 + UU_x - U(1 - U)] = 0 \quad 19$$

where $p \in [0,1]$ is an embedding parameter, u_0 represent the initial approximation that satisfies the solution of the equation

In the case of $p = 0$, then equation (19) becomes

$$H(U, 0) = U_t - u_0 = 0 \quad 20$$

In the case of $p = 1$, then equation (19) becomes

$$H(U, 1) = U_t + UU_x - U(1 - U) = 0 \quad 21$$

Now taking the Aboodh Transform on both sides of equation (19) we have

$$A[H(U, p)] = A[U_t - u_0 + p[u_0 + UU_x - U(1 - U)]] = 0 \quad 22$$

Using the initial property of Aboodh transform, we obtain

$$vU(x, v) - \frac{1}{v}U(x, 0) = A\{u_0 - p[u_0 + UU_x - U + U^2]\} \quad 23$$

$$U(x, v) = \frac{1}{v^2}U(x, 0) + \frac{1}{v}A\{u_0 - p[u_0 + UU_x - U + U^2]\} \quad 24$$

By applying the Inverse Aboodh transform on both sides of equation (24), we have

$$U(x, t) = U(x, 0) + A^{-1} \left\{ \frac{1}{v} A \{ u_0 - p [u_0 + UU_x - U + U^2] \} \right\} \quad 25$$

From the homotopy perturbation method, we use the embedding parameter p as a small parameter and assume that the solution of equation (25) can be written as power series of p as

$$U(x, t) = \sum_{n=0}^{\infty} p^n U_n(x, t) \quad 26$$

Substituting equation (26) into (25)

$$\sum_{n=0}^{\infty} p^n U_n(x, t) = U(x, 0) + A^{-1} \left\{ \frac{1}{v} A \{ u_0 - p [u_0 + \sum_{n=0}^{\infty} p^n \{ U_n U_{nx} - U_n + U_n^2 \}] \} \right\} \quad 27$$

Equating the coefficients of the corresponding powers of p in equation (27)

$$p^0 : U_0(x, t) = U(x, 0) + A^{-1} \left\{ \frac{1}{v} A \{ u_0 \} \right\}$$

$$U_0(x, t) = ae^{-x} + A^{-1} \left\{ \frac{1}{v} A \{ ae^{-x} \} \right\}$$

$$U_0(x, t) = ae^{-x} + ae^{-x}t$$

$$U_0(x, t) = ae^{-x}(1 + t)$$

$$p^1 : U_1(x, t) = -A^{-1} \left\{ \frac{1}{v} A \{ u_0 + U_0(U_0)_x - U_0 + U_0^2 \} \right\}$$

$$U_1(x, t) = -A^{-1} \left\{ \frac{1}{v} A \{ ae^{-x} + ae^{-x}(1 + t)(-ae^{-x}(1 + t)) - ae^{-x}(1 + t) + (ae^{-x}(1 + t))^2 \} \right\}$$

$$U_1(x, t) = \frac{1}{2!} ae^{-x}t^2$$

$$p^2 : U_2(x, t) = -A^{-1} \left\{ \frac{1}{v} A \{ U_0(U_1)_x + U_1(U_0)_x - U_1 + 2U_0U_1 \} \right\}$$

$$U_2(x, t) = -A^{-1} \left\{ \frac{1}{v} A \{ ae^{-x}(1 + t) \left(-\frac{1}{2!} ae^{-x}t^2 \right) + \frac{1}{2!} ae^{-x}t^2 (-ae^{-x}(1 + t)) - \frac{1}{2!} ae^{-x}t^2 + 2(ae^{-x}(1 + t)) \left(\frac{1}{2!} ae^{-x}t^2 \right) \} \right\}$$

$$U_2(x, t) = \frac{1}{3!} a e^{-x}t^3$$

$$p^3 : U_3(x, t) = -A^{-1} \left\{ \frac{1}{v} A \{ U_0(U_2)_x + U_1(U_1)_x + U_2(U_0)_x - U_2 + U_0U_1 + U_0U_2 + U_1U_1 + U_2U_0 \} \right\}$$

$$U_3(x, t) = -A^{-1} \left\{ \frac{1}{v} A \left\{ ae^{-x}(1 + t) \left(-\frac{1}{3!} ae^{-x}t^3 \right) + \frac{1}{2!} ae^{-x}t^2 \left(-\frac{1}{2!} ae^{-x}t^2 \right) + \frac{1}{3!} ae^{-x}t^3 (-ae^{-x}(1 + t)) - \frac{1}{3!} ae^{-x}t^3 + ae^{-x}(1 + t) \frac{1}{3!} ae^{-x}t^3 + \frac{1}{2!} ae^{-x}t^2 \left(\frac{1}{2!} ae^{-x}t^2 \right) + \frac{1}{3!} e^{-x}t^3 (ae^{-x}(1 + t)) \right\} \right\}$$

$$U_3(x, t) = \frac{1}{4!} a e^{-x}t^4$$

$$p^4 : U_4(x, t) = -A^{-1} \left\{ \frac{1}{v} A \{ U_0(U_3)_x + U_1(U_2)_x + U_2(U_1)_x + U_3(U_0)_x - U_3 + U_0U_3 + U_1U_2 + U_2U_1 + U_3U_0 \} \right\}$$

$$U_4(x, t) = -A^{-1} \left\{ \frac{1}{v} A \left\{ a e^{-x} (1+t) \left(\frac{1}{4!} a e^{-x} t^4 \right) + \frac{1}{2!} a e^{-x} t^2 \left(-\frac{1}{3!} a e^{-x} t^3 \right) \right. \right. \\ \left. \left. + \frac{1}{3!} a e^{-x} t^3 \left(-\frac{1}{2!} a e^{-x} t^2 \right) + \frac{1}{4!} a e^{-x} t^4 \left(-a e^{-x} (1+t) \right) - \frac{1}{4!} a e^{-x} t^4 \right\} \right. \\ \left. + a e^{-x} (1+t) \left(\frac{1}{4!} a e^{-x} t^4 \right) + a e^{-x} (1+t) \left(\frac{1}{3!} a e^{-x} t^3 \right) + \frac{1}{3!} a e^{-x} t^3 \left(\frac{1}{2!} a e^{-x} t^2 \right) \right. \\ \left. + \frac{1}{4!} a e^{-x} t^4 (a e^{-x} (1+t)) \right\}$$

$$U_4(x, t) = \frac{1}{5!} a e^{-x} t^5$$

⋮

$$p^m : U_m(x, t) = A^{-1} \left\{ -\frac{1}{v} A \left\{ \sum_{k=0}^{m-1} U_k (U_{m-k-1})_x - U_{m-1} + \sum_{k=0}^{m-1} U_k U_{m-k-1} \right\} \right\} \quad 28$$

The rest of the components of iteration formula can be obtained by following the same procedure.

Suppose the initial approximation has the form

$$U(x, 0) = u_0(x, t) = a e^{-x}$$

Thus, the exact solution can be obtained as follows:

$$u(x, t) = \lim_{p \rightarrow 1} U(x, t)$$

$$u(x, t) = U_0(x, t) + U_1(x, t) + U_2(x, t) + \dots$$

$$u(x, t) = a e^{-x} (1+t) + \frac{1}{2!} a e^{-x} t^2 + \frac{1}{3!} a e^{-x} t^3 + \frac{1}{4!} a e^{-x} t^4 + \frac{1}{5!} a e^{-x} t^5 + \dots \quad 29$$

$$u(x, t) = a e^{-x} (1+t+t^2+t^3+t^4+t^5+\dots) \quad 30$$

Hence, the solution can be written in the closed form as:

$$u(x, t) = a e^{t-x} \quad 31$$

Equation (31) is the exact solution for equation (17) which is the same as the solution obtained in (Aminikhah and Jamalain 2013).

Example 2

Consider the nonlinear homogenous partial differential equation (Jafari, Zabihi and Saidy 2008)

$$u_t + \frac{1}{2} (u^2)_x - u(1-u) = 0, \quad 0 \leq x \leq 1, \quad t > 0 \quad 32$$

with specified conditions

$$u(x, 0) = e^{-x} \quad 33$$

By applying the modified homotopy perturbation method, we construct the following homotopy:

$$H(U, p) = U_t - u_0 + p \left[u_0 + \frac{1}{2} U^2_x - U(1-U) \right] = 0 \quad 34$$

where $p \in [0,1]$ is an embedding parameter, u_0 represent the initial approximation that satisfies the solution of the equation

In the case of $p = 0$, then equation (34) becomes

$$H(U, 0) = U_t - u_0 = 0 \quad 35$$

In the case of $p = 1$, then equation (34) becomes

$$H(U, 1) = U_t + \frac{1}{2}U^2_x - U(1 - U) = 0 \tag{36}$$

Now taking the Aboodh Transform on both sides of equation (34) we have,

$$A[H(U, p)] = A \left[U_t - u_0 + p \left[u_0 + \frac{1}{2}U^2_x - U(1 - U) \right] \right] = 0 \tag{37}$$

Using the initial property of Aboodh transform, we obtain

$$vU(x, v) - \frac{1}{v}U(x, 0) = A \left\{ u_0 - p \left[u_0 + \frac{1}{2}U^2_x - U + U^2 \right] \right\} \tag{38}$$

$$U(x, v) = \frac{1}{v^2}U(x, 0) + \frac{1}{v}A \left\{ u_0 - p \left[u_0 + \frac{1}{2}U^2_x - U + U^2 \right] \right\} \tag{39}$$

By taking the Inverse Aboodh transform on both sides of equation (39), we have

$$U(x, t) = A^{-1} \left\{ \frac{1}{v^2}U(x, 0) \right\} + A^{-1} \left\{ \frac{1}{v}A \left\{ u_0 - p \left[u_0 + \frac{1}{2}U^2_x - U + U^2 \right] \right\} \right\} \tag{40}$$

$$U(x, t) = U(x, 0) + A^{-1} \left\{ \frac{1}{v}A \left\{ u_0 - p \left[u_0 + \frac{1}{2}U^2_x - U + U^2 \right] \right\} \right\} \tag{41}$$

From the homotopy perturbation method, we use the embedding parameter p as a small parameter and assume that the solution of equation (41) can be written as power series of p as:

$$U(x, t) = \sum_{n=0}^{\infty} p^n U_n(x, t) \tag{42}$$

Substituting equation (42) into (41)

$$\sum_{n=0}^{\infty} p^n U_n(x, t) = U(x, 0) + A^{-1} \left\{ \frac{1}{v}A \left\{ u_0 - p \left[u_0 + \sum_{n=0}^{\infty} p^n \left[\frac{1}{2}U^2_{nx} - U_n + U_n^2 \right] \right] \right\} \right\} \tag{43}$$

Equating the coefficients of the corresponding powers of p in equation (43)

$$p^0 : U_0(x, t) = U(x, 0) + A^{-1} \left\{ \frac{1}{v}A \{ u_0 \} \right\}$$

$$U_0(x, t) = e^{-x} + A^{-1} \left\{ \frac{1}{v}A \{ e^{-x} \} \right\}$$

$$U_0(x, t) = e^{-x} + e^{-x}t$$

$$U_0(x, t) = e^{-x}(1 + t)$$

$$p^1 : U_1(x, t) = -A^{-1} \left\{ \frac{1}{v}A \left\{ u_0 + \frac{1}{2}(U_0^2)_x - U_0 + U_0^2 \right\} \right\}$$

$$U_1(x, t) = -A^{-1} \left\{ \frac{1}{v}A \left\{ e^{-x} + \frac{1}{2}(-2ae^{-2x}(1+t)^2) - e^{-x}(1+t) + (e^{-2x}(1+t))^2 \right\} \right\}$$

$$U_1(x, t) = \frac{1}{2!}e^{-x}t^2$$

$$p^2 : U_2(x, t) = -A^{-1} \left\{ \frac{1}{v}A \{ U_0(U_1)_x + U_1(U_0)_x - U_1 + 2U_0U_1 \} \right\}$$

$$U_2(x, t) = -A^{-1} \left\{ \frac{1}{v}A \left\{ e^{-x}(1+t) \left(-\frac{1}{2!}e^{-x}t^2 \right) + \frac{1}{2!}e^{-x}t^2(-e^{-x}(1+t)) - \frac{1}{2!}e^{-x}t^2 + 2(e^{-x}(1+t)) \left(\frac{1}{2!}e^{-x}t^2 \right) \right\} \right\}$$

$$U_2(x, t) = \frac{1}{3!}e^{-x}t^3$$

$$p^3 : U_3(x, t) = -A^{-1} \left\{ \frac{1}{v}A \left\{ \frac{1}{2}(U_1^2)_x + U_0(U_2)_x + U_2(U_0)_x - U_2 + U_0U_2 + U_2U_0 + U_1^2 \right\} \right\}$$

$$\begin{aligned}
 U_3(x, t) &= -A^{-1} \left\{ \frac{1}{v} A \left(\frac{1}{2} \left(-\frac{1}{2} e^{-2x} t^4 \right) + e^{-x}(1+t) \left(-\frac{1}{3!} e^{-x} t^3 \right) + \frac{1}{3!} e^{-x} t^3 \left(-e^{-x}(1+t) \right) - \frac{1}{3!} e^{-x} t^3 \right. \right. \\
 &\quad \left. \left. + e^{-x}(1+t) \frac{1}{3!} e^{-x} t^3 + \frac{1}{3!} e^{-x} t^3 (e^{-x}(1+t)) + \frac{1}{4!} e^{-2x} t^4 \right) \right\} \\
 U_3(x, t) &= \frac{1}{4!} e^{-x} t^4 \\
 &\quad \vdots \\
 p^m : U_m(x, t) &= A^{-1} \left\{ -\frac{1}{v} A \left(\sum_{k=0}^{m-1} U_k (U_{m-k-1})_x - U_{m-1} + \sum_{k=0}^{m-1} U_k U_{m-k-1} \right) \right\} \tag{44}
 \end{aligned}$$

The rest of the components of iteration formula can be obtained by following the same procedure.

Suppose the initial approximation has the form

$$U(x, 0) = u_0(x, t) = e^{-x}$$

Thus, the exact solution can be obtained as follows:

$$u(x, t) = \lim_{p \rightarrow 1} U(x, t)$$

$$u(x, t) = U_0(x, t) + U_1(x, t) + U_2(x, t) + \dots$$

$$u(x, t) = e^{-x}(1+t) + \frac{1}{2!} e^{-x} t^2 + \frac{1}{3!} e^{-x} t^3 + \frac{1}{4!} e^{-x} t^4 + \dots \tag{45}$$

$$u(x, t) = e^{-x}(1+t+t^2+t^3+t^4+\dots) \tag{46}$$

Hence, the solution can be written in the closed form as:

$$u(x, t) = e^{t-x} \tag{47}$$

Equation (47) is the exact solution for equation (32) which is the same as the solution obtained in (Jafari, Zabihi and Saidy 2008).

Example 3

Consider the following non homogeneous nonlinear gas dynamic equation (Aminikhah and Jamalain 2013)

$$u_t + uu_x - u(1-u) = -e^{t-x} \tag{48}$$

subject to initial condition

$$u(x, 0) = 1 - e^{-x} \tag{49}$$

By applying the modified homotopy perturbation method, we construct the following homotopy:

$$H(U, p) = U_t - u_0 + p[u_0 + UU_x - U(1-U) + e^{t-x}] = 0 \tag{50}$$

where $p \in [0,1]$ is an embedding parameter, u_0 represent the initial approximation that satisfies the solution of the equation

In the case of $p = 0$, then equation (50) becomes

$$H(U, 0) = U_t - u_0 = 0 \tag{51}$$

In the case of $p = 1$, then equation (50) becomes

$$H(U, 1) = U_t + UU_x - U(1-U) + e^{t-x} = 0 \tag{52}$$

Now taking the Aboodh Transform on both sides of equation (50) we have

$$A[H(U, p)] = A[U_t - u_0 + p[u_0 + UU_x - U(1 - U) + e^{t-x}]] = 0 \tag{53}$$

Using the initial property of Aboodh transform, we obtain

$$vU(x, v) - \frac{1}{v}U(x, 0) = A\{u_0 - p[u_0 + UU_x - U + U^2 + e^{t-x}]\} \tag{54}$$

$$U(x, v) = \frac{1}{v^2}U(x, 0) + \frac{1}{v}A\{u_0 - p[u_0 + UU_x - U + U^2 + e^{t-x}]\} \tag{55}$$

By taking the Inverse Aboodh transform on both sides of equation (55), we have

$$U(x, t) = A^{-1}\left\{\frac{1}{v^2}U(x, 0)\right\} + A^{-1}\left\{\frac{1}{v}A\{u_0 - p[u_0 + UU_x - U + U^2 + e^{t-x}]\}\right\} \tag{56}$$

$$U(x, t) = U(x, 0) + A^{-1}\left\{\frac{1}{v}A\{u_0 - p[u_0 + UU_x - U + U^2 + e^{t-x}]\}\right\} \tag{57}$$

From the homotopy perturbation method, we use the embedding parameter p as a small parameter and assume that the solution of equation (57) can be written as power series of p as

$$U(x, t) = \sum_{n=0}^{\infty} p^n U_n(x, t) \tag{58}$$

Substituting equation (58) into (57)

$$\sum_{n=0}^{\infty} p^n U_n(x, t) = U(x, 0) + A^{-1}\left\{\frac{1}{v}A\{u_0 - p[u_0 + \sum_{n=0}^{\infty} p^n \{U_n U_{nx} - U_n + U_n^2\} + e^{t-x}]\}\right\} \tag{59}$$

Equating the coefficients of the corresponding powers of p in equation (59)

$$p^0 : U_0(x, t) = U(x, 0) + A^{-1}\left\{\frac{1}{v}A\{u_0\}\right\}$$

$$U_0(x, t) = 1 - e^{-x} + A^{-1}\left\{\frac{1}{v}A\{1 - e^{-x}\}\right\}$$

$$U_0(x, t) = 1 - e^{-x} + t - e^{-x}t$$

$$U_0(x, t) = 1 + t - e^{-x}(1 + t)$$

$$p^1 : U_1(x, t) = -A^{-1}\left\{\frac{1}{v}A\{u_0(x, t) + U_0(U_0)_x - U_0 + U_0^2 + e^{t-x}\}\right\}$$

$$U_1(x, t) = -A^{-1}\left\{\frac{1}{v}A\left\{(1 - e^{-x}) + (1 + t - e^{-x}(1 + t))(e^{-x}(1 + t)) - (1 + t - e^{-x}(1 + t)) + (1 + t - e^{-x}(1 + t))^2 + e^{t-x}\right\}\right\}$$

$$U_1(x, t) = -t - \frac{t^2}{2} - \frac{t^3}{3} - e^{t-x} + \frac{1}{6}e^{-x}(6 + 6t + 3t^2 + 2t^3)$$

$$p^2 : U_2(x, t) = -A^{-1}\left\{\frac{1}{v}A\{U_0(U_1)_x + U_1(U_0)_x - U_1 + 2U_0U_1\}\right\}$$

$$U_2(x, t) = -A^{-1} \left\{ \frac{1}{v} A \left\{ (1+t - e^{-x}(1+t)) \left(e^{t-x} - \frac{1}{6} e^{-x}(6+6t+3t^2+2t^3) \right) \right. \right. \\ \left. \left. + \left(-t - \frac{t^2}{2} - \frac{t^3}{3} - e^{t-x} + \frac{1}{6} e^{-x}(6+6t+3t^2+2t^3) \right) (e^{-x}(1+t)) \right. \right. \\ \left. \left. - \left(-t - \frac{t^2}{2} - \frac{t^3}{3} - e^{t-x} + \frac{1}{6} e^{-x}(6+6t+3t^2+2t^3) \right) \right. \right. \\ \left. \left. + 2(1+t - e^{-x}(1+t)) \left(-t - \frac{t^2}{2} - \frac{t^3}{3} - e^{t-x} + \frac{1}{6} e^{-x}(6+6t+3t^2+2t^3) \right) \right\} \right\}$$

$$U_2(x, t) = \frac{t^2}{2} + \frac{5}{6}t^3 + \frac{1}{3}t^4 + \frac{2}{15}t^5 - \frac{1}{30}e^{-x}(-30 + 30t^2 + 25t^3 + 10t^4 + 4t^5) + e^{t-x} (t - 1)$$

$$p^3 : U_3(x, t) = -A^{-1} \left\{ \frac{1}{v} A \{ U_0(U_2)_x + U_1(U_1)_x + U_2(U_0)_x - U_2 + U_0U_1 + U_0U_2 + U_1^2 + U_2U_0 \} \right\}$$

$$U_3(x, t) = -A^{-1} \left\{ \frac{1}{v} A \left\{ (1+t - e^{-x}(1+t)) \left(\frac{1}{30} e^{-x}(-30 + 30t^2 + 25t^3 + 10t^4 + 4t^5) - e^{t-x} (t - 1) \right) \right. \right. \\ \left. \left. + \left(-t - \frac{t^2}{2} - \frac{t^3}{3} - e^{t-x} + \frac{1}{6} e^{-x}(6+6t+3t^2+2t^3) \right) \left(e^{t-x} \right. \right. \right. \\ \left. \left. - \frac{1}{6} e^{-x}(6+6t+3t^2+2t^3) \right) \right. \right. \\ \left. \left. + \left(\frac{t^2}{2} + \frac{5}{6}t^3 + \frac{1}{3}t^4 + \frac{2}{15}t^5 - \frac{1}{30} e^{-x}(-30 + 30t^2 + 25t^3 + 10t^4 + 4t^5) \right. \right. \right. \\ \left. \left. + e^{t-x} (t - 1) \right) (e^{-x}(1+t)) \right. \right. \\ \left. \left. - \left(\frac{t^2}{2} + \frac{5}{6}t^3 + \frac{1}{3}t^4 + \frac{2}{15}t^5 - \frac{1}{30} e^{-x}(-30 + 30t^2 + 25t^3 + 10t^4 + 4t^5) + e^{t-x} (t - 1) \right) \right. \right. \\ \left. \left. + (1+t - e^{-x}(1+t)) \left(-t - \frac{t^2}{2} - \frac{t^3}{3} - e^{t-x} + \frac{1}{6} e^{-x}(6+6t+3t^2+2t^3) \right) \right. \right. \\ \left. \left. + (1+t - e^{-x}(1+t)) \left(\frac{t^2}{2} + \frac{5}{6}t^3 + \frac{1}{3}t^4 + \frac{2}{15}t^5 \right. \right. \right. \\ \left. \left. - \frac{1}{30} e^{-x}(-30 + 30t^2 + 25t^3 + 10t^4 + 4t^5) + e^{t-x} (t - 1) \right) \right. \right. \\ \left. \left. + \left(-t - \frac{t^2}{2} - \frac{t^3}{3} - e^{t-x} + \frac{1}{6} e^{-x}(6+6t+3t^2+2t^3) \right)^2 \right. \right. \\ \left. \left. + \left(\frac{t^2}{2} + \frac{5}{6}t^3 + \frac{1}{3}t^4 + \frac{2}{15}t^5 - \frac{1}{30} e^{-x}(-30 + 30t^2 + 25t^3 + 10t^4 + 4t^5) \right. \right. \right. \\ \left. \left. + e^{t-x} (t - 1) \right) (1+t - e^{-x}(1+t)) \right\} \right\}$$

$$U_3(x, t) = -\frac{1}{2}t^3 - \frac{17}{24}t^4 - \frac{7}{12}t^5 - \frac{17}{90}t^6 - \frac{17}{315}t^7 - \frac{1}{6}e^{t-x}(6 - 6t + 3t^2 + 2t^3) \\ + \frac{1}{1260}e^{-x} (1260 + 840t^3 + 1155t^4 + 735t^5 + 238t^6 \\ + 68t^7)$$

$$p^4 : U_4(x, t) = -A^{-1} \left\{ \frac{1}{v} A \{ U_0(U_3)_x + U_1(U_2)_x + U_2(U_1)_x + U_3(U_0)_x - U_3 + U_0U_3 + 2U_1U_2 + U_3U_0 \} \right\}$$

$$\begin{aligned}
 U_4(x, t) = & -A^{-1} \left\{ \frac{1}{v} A \left\{ (1+t - e^{-x}(1+t)) \left(\frac{1}{6} e^{t-x} (6 - 6t + 3t^2 + 2t^3) \right. \right. \right. \\
 & - \frac{1}{1260} e^{-x} (1260 + 840t^3 + 1155t^4 + 735t^5 + 238t^6 + 68t^7) \left. \right. \left. \right) \\
 & + \left(-t - \frac{t^2}{2} - \frac{t^3}{3} - e^{t-x} \right. \\
 & + \left. \frac{1}{6} e^{-x} (6 + 6t + 3t^2 + 2t^3) \right) \left(\frac{1}{30} e^{-x} (-30 + 30t^2 + 25t^3 + 10t^4 + 4t^5) \right. \\
 & - \left. e^{t-x} (t-1) \right) \\
 & + \left(\frac{t^2}{2} + \frac{5}{6} t^3 + \frac{1}{3} t^4 + \frac{2}{15} t^5 - \frac{1}{30} e^{-x} (-30 + 30t^2 + 25t^3 + 10t^4 + 4t^5) \right. \\
 & + \left. e^{t-x} (t-1) \right) \left(e^{t-x} - \frac{1}{6} e^{-x} (6 + 6t + 3t^2 + 2t^3) \right) \\
 & + \left(-\frac{1}{2} t^3 - \frac{17}{24} t^4 - \frac{7}{12} t^5 - \frac{17}{90} t^6 - \frac{17}{315} t^7 - \frac{1}{6} e^{t-x} (6 - 6t + 3t^2 + 2t^3) \right. \\
 & + \left. \frac{1}{1260} e^{-x} (1260 + 840t^3 + 1155t^4 + 735t^5 + 238t^6 + 68t^7) \right) (e^{-x}(1+t)) \\
 & - \left(-\frac{1}{2} t^3 - \frac{17}{24} t^4 - \frac{7}{12} t^5 - \frac{17}{90} t^6 - \frac{17}{315} t^7 - \frac{1}{6} e^{t-x} (6 - 6t + 3t^2 + 2t^3) \right. \\
 & + \left. \frac{1}{1260} e^{-x} (1260 + 840t^3 + 1155t^4 + 735t^5 + 238t^6 + 68t^7) \right) \\
 & + (1+t - e^{-x}(1+t)) \left(-\frac{1}{2} t^3 - \frac{17}{24} t^4 - \frac{7}{12} t^5 - \frac{17}{90} t^6 - \frac{17}{315} t^7 \right. \\
 & - \left. \frac{1}{6} e^{t-x} (6 - 6t + 3t^2 + 2t^3) \right) \\
 & + \left. \frac{1}{1260} e^{-x} (1260 + 840t^3 + 1155t^4 + 735t^5 + 238t^6 + 68t^7) \right) \\
 & + 2 \left(-t - \frac{t^2}{2} - \frac{t^3}{3} - e^{t-x} + \frac{1}{6} e^{-x} (6 + 6t + 3t^2 + 2t^3) \right) \left(\frac{t^2}{2} + \frac{5}{6} t^3 + \frac{1}{3} t^4 + \frac{2}{15} t^5 \right. \\
 & - \left. \frac{1}{30} e^{-x} (-30 + 30t^2 + 25t^3 + 10t^4 + 4t^5) + e^{t-x} (t-1) \right) \\
 & + \left(-\frac{1}{2} t^3 - \frac{17}{24} t^4 - \frac{7}{12} t^5 - \frac{17}{90} t^6 - \frac{17}{315} t^7 - \frac{1}{6} e^{t-x} (6 - 6t + 3t^2 + 2t^3) \right. \\
 & + \left. \frac{1}{1260} e^{-x} (1260 + 840t^3 + 1155t^4 + 735t^5 + 238t^6 + 68t^7) \right) (1+t \\
 & - e^{-x}(1+t)) \left. \right\} \left. \right\}
 \end{aligned}$$

$$\begin{aligned}
 U_4(x, t) = & \frac{3}{8} t^4 + \frac{31}{40} t^5 + \frac{23}{36} t^6 + \frac{113}{315} t^7 + \frac{31}{315} t^8 + \frac{62}{2835} t^8 \\
 & + \frac{1}{45360} e^{-x} (-45360 + 28350t^4 + 41958t^5 + 32823t^6 + 16272t^7 + 4464t^8 + 992t^9) \\
 & + \frac{1}{30} e^{t-x} (-30 + 30t - 15t^2 + 5t^3 + 10t^4 + 4t^5)
 \end{aligned}$$

⋮

$$p^m : U_m(x, t) = A^{-1} \left\{ -\frac{1}{v} A \left\{ \sum_{k=0}^{m-1} U_k (U_{m-k-1})_x - U_{m-1} + \sum_{k=0}^{m-1} U_k U_{m-k-1} \right\} \right\} \quad 60$$

The rest of the components of iteration formula can be obtained by following the same procedure.

Suppose the initial approximation has the form

$$U(x, 0) = u_0(x, t) = 1 - e^{-x}$$

Thus, the exact solution can be obtained as follows:

$$u(x, t) = \lim_{p \rightarrow 1} U(x, t)$$

$$u(x, t) = U_0(x, t) + U_1(x, t) + U_2(x, t) + U_3(x, t) \dots$$

$$u(x, t) = (1 + t - e^{-x}(1 + t)) + \left(-t - \frac{t^2}{2} - \frac{t^3}{3} - e^{-x} + \frac{1}{6} e^{-x}(6 + 6t + 3t^2 + 2t^3) \right) + \left(\frac{t^2}{2} + \frac{5}{6} t^3 + \frac{1}{3} t^4 + \frac{2}{15} t^5 - \frac{1}{30} e^{-x}(-30 + 30t^2 + 25t^3 + 10t^4 + 4t^5) + e^{-x} (t - 1) \right) + \left(-\frac{1}{2} t^3 - \frac{17}{24} t^4 - \frac{7}{12} t^5 - \frac{17}{90} t^6 - \frac{17}{315} t^7 - \frac{1}{6} e^{-x}(6 - 6t + 3t^2 + 2t^3) + \frac{1}{1260} e^{-x} (1260 + 840t^3 + 1155t^4 + 735t^5 + 238t^6 + 68t^7) \right) + \left(\frac{3}{8} t^4 + \frac{31}{40} t^5 + \frac{23}{36} t^6 + \frac{113}{315} t^7 + \frac{31}{315} t^8 + \frac{62}{2835} t^8 + \frac{1}{45360} e^{-x} (-45360 + 28350t^4 + 41958t^5 + 32823t^6 + 16272t^7 + 4464t^8 + 992t^9) + \frac{1}{30} e^{-x}(-30 + 30t - 15t^2 + 5t^3 + 10t^4 + 4t^5) \right) + \dots \quad 61$$

$$u(x, t) \cong 1 + 4e^{-x} - 5e^{-x}t - \frac{13}{35}t^2 - \frac{172}{945}t^3 - \frac{11}{60}e^{-x}t^4 + \frac{7}{24}e^{-x}t^5 + \frac{16361}{40320}e^{-x}t^6 + \frac{172}{945}e^{-x}t^7 - \frac{17}{90}e^{-x}t^8 - \frac{17}{315}e^{-x}t^9 + \frac{1382}{155925}e^{-x}t^{10} + \frac{691}{14175}e^{-x}t^{11} - \frac{21}{80}t^6 - \frac{79}{168}t^7 - \frac{3}{2}e^{-x}t^2 + 4e^{-x}t + \frac{407}{720}e^{-x}t^7 - \frac{1}{24}e^{-x}t^4 + \frac{33}{80}e^{-x}t^6 + \frac{1}{120}e^{-x}t^5 + \frac{1}{6}e^{-x}t^3 - \frac{691}{14175}t^{10} - \frac{1382}{155925}t^{11} - \frac{1}{2}e^{-x}t^2 \quad 62$$

Equation (62) is the exact solution for equation (48) which is the same as the solution obtained in (Aminikhah and Jamalian 2013).

4. CONCLUSION

In this paper, we introduced the Modified Aboodh Homotopy Perturbation Method which is the combination of Aboodh transform and Modified homotopy perturbation method to solve nonlinear gas dynamic equations. The main advantage of this method is that, it provides the user an analytical approximation to the solution in series of rapidly convergent sequence with elegantly computed terms. The results obtained show that the method is trustworthy and introduces a significant advancement in solving nonlinear partial differential equations over existing methods.

REFERENCE

Aboodh, K. S. (2013) "The New Integral Transform Aboodh Transform", *Global Journal of Pure and Applied Mathematics* ISSN0973-1768, vol. 9, no. 1, pp. 35-43.
 Aboodh, K. S. (2014), "Application of New "Aboodh Transform" to Partial Differential Equations", *Global Journal of*

pure and Applied Mathematics, vol. 10, no. 2, pp. 249-254.

Aboodh, K. S. (2015), Homotopy perturbation method and Aboodh Transform for solving nonlinear partial differential equations, *Journal of pure and applied Mathematics*, Vol.4, no. 5, pp. 219-224.
 Ames W. F. (1965), *Nonlinear Partial Differential Equations in Engineering*, Academic Press, New York, NY, USA.
 Aminikhah H. and Jamalian A. (2013), "Numerical approximation for nonlinear gas dynamic equation" *The International Journal of Partial Differential Equations* Volume 2013, Article ID 846749, 7 pages.
 Bellman R. (1964), *Perturbation Techniques in Mathematics, Physics and Engineering*, Holt, Rinehart and Winston, New York.
 Cole J.D. (1968), *Perturbation Methods in Applied Mathematics*, Blaisedell, Waltham, MA.
 Elizarova T. G., (2009), *Quasi-Gas Dynamic Equations*, Computational Fluid and Solid Mechanics, Springer, Dordrecht, Netherlands.

- Evans D. J. and Bulut H., (2002) "A new approach to the gas dynamics equation: an application of the decomposition method," *International Journal of Computer Mathematics*, vol. 79, no. 7, pp.817– 822.
- He, J.H. (1999), Homotopy perturbation technique, *Computer Methods in Applied Mechanics and Engineering*, vol. 178, no. 3-4, pp. 257–262, 1999.
- He, J.H. (2000), Coupling method of a homotopy technique and a perturbation technique for non-linear problems, *International Journal of Non-Linear Mechanics*, vol. 35, no.1, pp. 37–43.
- He, J.H. (2000) "Application of homotopy perturbation method to nonlinear wave equations," *Chaos, Solitons and Fractals*, vol. 26, no. 3, pp. 695–700, 2005.
- He. J. H. (2003), Homotopy Perturbation Method: A new nonlinear analytical technique. *Applied Mathematics and Computation*, vol. 135, pp 73-79.
- He, J.H. (2004), "The homotopy perturbation method for nonlinear oscillators with discontinuities," *Applied Mathematics and Computation*, vol. 151, no. 1, pp. 287–292.
- He, J.H. (2005) "Limit cycle and bifurcation of nonlinear problems" *Chaos, Solitons and Fractals*, vol. 26, no. 3, pp. 827–833.
- He, J.H. (2006), "Addendum: new interpretation of homotopy perturbation method," *International Journal of Modern Physics B*, vol. 20, no. 18, pp. 2561–2568.
- He. J. H. (2008), Recent developments of the homotopy perturbation method. *Topological Methods in Nonlinear Analysis*, vol. 31, no.2, pp 205-209.
- Jafari H., Zabihi and Saidy M. (2008) "Application of Homotopy –Perturbation method for solving gas dynamics equation" ,vol. 2, no.48, pp. 2393-2396.
- Jafari, H. Chun C., Seifi, S. and Saeidy M., (2009) "Analytical solution for nonlinear gas dynamic equation by homotopy analysis method," *Applications and Applied Mathematics*, vol. 4, no. 1, pp. 149–154.
- Mohand M. Abdelrahim Mahgoub (2016), "Homotopy Perturbation Method and Aboodh Transform for solving sine-Gorden and Klein – Gorden Equations" *International Journal of Engineering Sciences & Research Technology* Vol.5, no. 10.
- Polyanin A. D. and Zaitsev V. F., (2004) *Handbook of Nonlinear Partial Differential Equations*, Chapman & Hall/CRC, Boca Raton, Fla, USA,.
- Rasulov M. and Karaguler T.,(2003), "Finite difference schemes for solving system equations of gas dynamic in a class of discontinuous functions," *Applied Mathematics and Computation*, vol.143, no. 1, pp. 145–164, 2003.
- Sahoo M. (2015), Some applications of homotopy perturbation method, M.Sc Thesis, National Institute of Technology, Rourkela, pp1-25.